

BEFORE TO AFTER PHOTOS

Before treatment
(all photos unretouched)

Session 2 results

Session 3 results

Month 7 after first injection results



Indication:

Sculptra®Aesthetic is indicated for the correction of shallow to deep nasolabial fold (smile lines) contour deficiencies and other facial wrinkles, which are treated with the appropriate injection technique in healthy people.

Important Safety Information:

You should not use Sculptra Aesthetic if you are allergic to any ingredient of the product or have a history of keloid formation or hypertrophic scarring. Sculptra Aesthetic should not be injected while you have an active skin infection or inflammation in the treatment area and should not be injected into the red area of the lip. Use in the skin near the eyes is not recommended.

Side effects of Sculptra Aesthetic may include injection site discomfort, redness, bruising, bleeding, itching and swelling. Other side effects may include small lumps under the skin that are sometimes noticeable when pressing on the treated area. Larger lumps, some with delayed onset with or without inflammation or skin discoloration, have also been reported. In a key clinical study the numbers of small and larger lumps were low and most resolved without treatment.

For more information, including full prescribing information, please visit www.SculptraAesthetic.com or www.sanofi-aventis.com.

Sculptra®Aesthetic is not for use in the lips. Safety and effectiveness of Sculptra®Aesthetic has not been evaluated in areas other than nasolabial wrinkles and folds in clinical trials. Sculptra®Aesthetic is not recommended for use in the periorbital area.

On average, 3 treatment sessions are needed over a period of a few months to produce gradual, subtle results that can last up to two years.

Please see full prescribing information at www.SculptraAesthetic.com.

sculptra®aesthetic
injectable poly-L-lactic acid

US.POL.09.07.014